

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Vibramycin Capsules 100 mg

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

100 mg doxycycline as doxycycline hyclate.

Excipients with known effect:

Each capsule contains 104 mg lactose (as monohydrate).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Capsule, hard.

Dark green opaque capsules, printed "Pfizer Vibracina" in off-white colour.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Vibramycin has been found clinically effective in the treatment of a variety of infections caused by susceptible strains of Gram-positive and Gram-negative bacteria and certain other micro-organisms.

Respiratory tract infections:

Pneumonia and other lower respiratory tract infections due to susceptible strains of *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Klebsiella pneumoniae*, *Moraxella catarrhalis* and other organisms. *Mycoplasma pneumoniae*. Treatment of chronic bronchitis, sinusitis.

Urinary tract infections:

Infections caused by susceptible strains of *Klebsiella* species, *Enterobacter* species, *Escherichia coli*, *Streptococcus faecalis* and other organisms.

Sexually transmitted diseases:

Infections due to *Chlamydia trachomatis* including uncomplicated urethral, endocervical or rectal infections. Non-gonococcal urethritis caused by *Ureaplasma urealyticum* (T-mycoplasma). Vibramycin is also indicated in infections due to *Calymmatobacterium granulomatis*. Vibramycin is an alternative drug in the treatment of gonorrhoea and syphilis.

Since Vibramycin is a member of the tetracycline series of antibiotics, it may be expected to be useful in the treatment of infections which respond to other tetracyclines such as:

Ophthalmic infections:

Vibramycin is indicated in the treatment of trachoma, although the infectious agent is not always eliminated, as judged by immunofluorescence. Inclusion conjunctivitis may be treated with oral Vibramycin alone or in combination with topical agents.

Rickettsial infections:

Rocky Mountain spotted fever, typhus group, Q fever and Coxiella endocarditis.

Other infections:

Psittacosis, brucellosis (in combination with streptomycin), cholera, bubonic plague, louse and tick-borne relapsing fever including stage 1 and stage 2 Lyme disease, leptospirosis, tularaemia glanders, chloroquine-resistant falciparum malaria and acute intestinal amoebiasis (as an adjunct to amoebicides). Infections due to susceptible strains of *Bacteroides* species, *Listeria* species and *Bacillus anthracis*.

Vibramycin is an alternative drug in the treatment of leptospirosis, gas gangrene and tetanus.

Vibramycin is indicated for prophylaxis in the following conditions: Scrub typhus, travellers' diarrhoea (enterotoxigenic *Escherichia coli*), leptospirosis, malaria and cholera.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

Adults and children aged 12 years to less than 18 years

The usual dose of Vibramycin for the treatment of acute infections in adults and children aged 12 years to less than 18 years is 200 mg on the first day (administered as a single dose or divided into two equal doses with a 12 hour interval), followed by a maintenance dose of 100 mg/day. In the management of more severe infections (particularly chronic infections of the urinary tract) 200 mg daily should be given throughout the treatment period.

Children aged 8 years to less than 12 years (see section 4.4)

The use of doxycycline for the treatment of acute infections in children aged 8 years to less than 12 years should be carefully justified in situations where other drugs are not available, are not likely to be effective or are contraindicated.

In such circumstance, the doses for the treatment of acute infections are:

- For children 45 kg or less - Initial dose: 4.4 mg/kg (in single or 2 divided doses) with maintenance dose: 2.2 mg/kg (in single or 2 divided doses). In the management of more severe infections, up to 4.4 mg/kg should be given throughout treatment.
- For children over 45 kg - Dose administered for adults should be used.

Children aged from birth to less than 8 years

Doxycycline should not be used in children aged younger than 8 years due to the risk of teeth discolouration (section 4.4 and 4.8)

Method of administration

Vibramycin capsules should be taken with adequate amounts of fluid (at least 100mls of water). This should be done in the sitting or standing position and the patient should be

advised to remain upright for at least thirty minutes after taking a dose. Vibramycin capsules should be taken well before bedtime to reduce the risk of oesophageal irritation and ulceration. If gastric irritation occurs, it is recommended that Vibramycin be given with food or milk. Studies indicate that the absorption of Vibramycin is not notably influenced by simultaneous ingestion of food or milk.

Exceeding the recommended dosage may result in an increased incidence of side effects. Therapy should be continued at least 24 to 48 hours after symptoms and fever have subsided.

When used in streptococcal infections, therapy should be continued for 10 days to prevent the development of rheumatic fever or glomerulonephritis.

Dosage recommendations in specific infections:

Sexually transmitted diseases 100 mg twice daily for 7 days is recommended in the following infections: uncomplicated urethral, endocervical or rectal infection caused by *Chlamydia trachomatis*, non-gonococcal urethritis caused by *Ureaplasma urealyticum*.

Uncomplicated gonococcal infections (except anorectal infections in men) Doxycycline 100 mg twice daily for 7 days together with intramuscular ceftriaxone.

Acute epididymo-orchitis caused by Chlamydia trachomatis or Neisseria gonorrhoea
Doxycycline 100 mg twice daily for 10 days together with intramuscular ceftriaxone.

Primary and secondary syphilis Non-pregnant penicillin-allergic patients who have primary or secondary syphilis can be treated with the following regimen: doxycycline 100 mg orally twice daily for two weeks, as an alternative to penicillin therapy.

Louse and tick-borne relapsing fevers and louse borne typhus A single dose of 100 to 200 mg according to severity.

Early Lyme disease (Stage 1 and 2) 100 mg twice daily for 10-30 days according to clinical signs, symptoms and response.

Chloroquine-resistant falciparum malaria 200 mg daily for at least 7 days. Due to the potential severity of the infection, a rapid-acting schizonticide such as quinine should always be given in conjunction with Vibramycin; quinine dosage recommendations vary in different areas.

Prophylaxis of malaria 100 mg daily in adults. Prophylaxis can begin 1-2 days before travel to malarious areas. It should be continued daily during travel in the malarious areas and for 4 weeks after the traveller leaves the malarious area.

For the treatment and selective prophylaxis of cholera in adults 300 mg as a single dose.

For the prevention of scrub typhus 200 mg as a single dose, once weekly.

For the prevention of travellers' diarrhoea in adults 200 mg on the first day of travel (administered as a single dose or as 100 mg every 12 hours) followed by 100 mg daily throughout the stay in the area. Data on the use of the drug prophylactically are not available beyond 21 days.

For the treatment of leptospirosis 100 mg twice daily for 7 days.

For the prevention of leptospirosis 200 mg once each week throughout the stay in the area and 200 mg at the completion of the trip. Data on the use of the drug prophylactically are not available beyond 21 days.

Infections due to susceptible strains of Bacillus anthracis

Adults: 100 mg of doxycycline, by mouth, twice a day for 60 days.

Paediatric population See sections 4.3 and 4.4.

Use in the elderly Vibramycin may be prescribed in the usual dose with no special precautions. No dosage adjustment is necessary in the presence of renal impairment.

Use in patients with impaired hepatic function Vibramycin should be administered with caution to patients with hepatic impairment or those receiving potentially hepatotoxic drugs.

Use in patients with renal impairment Studies to date have indicated that administration of Vibramycin at the usual recommended doses does not lead to accumulation of the antibiotic in patients with renal impairment.

Rocky Mountain spotted fever

Vibramycin is the first line treatment for adults and children of all ages:

ADULTS: 100 mg every 12 hours.

CHILDREN: weighing less than 45 kg: 2.2 mg/kg body weight given twice a day. Children weighing 45 kg or more should receive the adult dose (see section 4.4 paediatric population).

Patients should be treated for at least 3 days after the fever subsides and until there is evidence of clinical improvement. Minimum course of treatment is 5-7 days.

4.3 Contraindications

Hypersensitivity to doxycycline, or to any other tetracyclines or to any of the excipients listed in section 6.1.

Obstructive oesophageal disorders, such as stricture or achalasia.

Pregnancy Vibramycin is contraindicated in pregnancy. It appears that the risks associated with the use of tetracyclines during pregnancy are predominantly due to effects on teeth and skeletal development. (see section 4.4 regarding use during tooth development).

Nursing Mothers Tetracyclines are excreted into milk and are therefore contraindicated in nursing mothers. (see section 4.4 regarding use during tooth development).

Paediatric population Vibramycin is contraindicated in children under the age of 8 years. As with other tetracyclines, Vibramycin forms a stable calcium complex in any bone-forming tissue. A decrease in the fibula growth rate has been observed in prematures given oral tetracyclines in doses of 25 mg/kg every 6 hours. This reaction was shown to be reversible when the drug was discontinued. (See above about use during tooth development).

4.4 Special warnings and precautions for use

Paediatric population The use of drugs of the tetracycline class during tooth development (last half of pregnancy; infancy and childhood to the age of 8 years) may cause permanent discolouration of the teeth (yellow-grey-brown). This adverse reaction is more common during long-term use of the drugs but has been observed following repeated short-term courses. Enamel hypoplasia has also been reported. Use doxycycline in paediatric patients aged younger than 8 years only when the potential benefits are expected to outweigh the risks in severe or life-threatening conditions (e.g. Rocky Mountain spotted fever), only when there are no adequate alternative therapies.

Although the risk of permanent teeth staining is rare in children aged 8 years to less than 12 years, the use of doxycycline should be carefully justified in situations where other drugs are not available, are not likely to be effective or are contraindicated.

Use in Patients with Impaired Hepatic Function Vibramycin should be administered with caution to patients with hepatic impairment or those receiving potentially hepatotoxic drugs.

Abnormal hepatic function has been reported rarely and has been caused by both the oral and parenteral administration of tetracyclines, including doxycycline.

Use in Patients with Renal Impairment Excretion of doxycycline by the kidney is about 40%/72 hours in individuals with normal renal function. This percentage excretion may fall to a range as low as 1-5%/72 hours in individuals with severe renal insufficiency (creatinine clearance below 10ml/min). Studies have shown no significant difference in the serum half-life of doxycycline in individuals with normal and severely impaired renal function. Haemodialysis does not alter the serum half-life of doxycycline. The anti-anabolic action of the tetracyclines may cause an increase in blood urea. Studies to date indicate that this anti-anabolic effect does not occur with the use of Vibramycin in patients with impaired renal function.

General Severe skin reactions, such as exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in patients receiving doxycycline (see section 4.8). If severe skin reactions occur, doxycycline should be discontinued immediately and appropriate therapy should be instituted.

Photosensitivity Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines, including doxycycline. Patients likely to be exposed to direct sunlight or ultraviolet light should be advised that this reaction can occur with tetracycline drugs and treatment should be discontinued at the first evidence of skin erythema.

Microbiological overgrowth The use of antibiotics may occasionally result in over-growth of non susceptible organisms. Constant observation of the patient is essential. If a resistant organism appears, the antibiotic should be discontinued and appropriate therapy instituted.

General Benign intracranial hypertension (pseudotumor cerebri) has been associated with the use of tetracyclines including doxycycline. Benign intracranial hypertension (pseudotumor cerebri) is usually transient, however cases of permanent visual loss secondary to benign intracranial hypertension (pseudotumor cerebri) have been reported with tetracyclines including doxycycline. If visual disturbance occurs during treatment, prompt ophthalmologic evaluation is warranted. Since intracranial pressure can remain elevated for weeks after drug

cessation patients should be monitored until they stabilize. Concomitant use of isotretinoin and doxycycline should be avoided because isotretinoin is also known to cause benign intracranial hypertension (pseudotumor cerebri).

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including doxycycline, and has ranged in severity from mild to life-threatening. It is important to consider this diagnosis in patients who present with diarrhoea subsequent to the administration of antibacterial agents.

Clostridium difficile associated diarrhoea (CDAD) has been reported with use of nearly all antibacterial agents, including doxycycline, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhoea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

Oesophagitis Cases of oesophageal injuries (oesophagitis and ulceration), sometimes serious, have been reported. Patients should be instructed to take doxycycline capsules with plenty of water (at least 100ml), remain upright and not take their treatment before going to bed (see section 4.2). Withdrawal of doxycycline and investigation of oesophageal disorder should be considered if symptoms such as dyspepsia or retrosternal pain occur. Caution is required in the treatment of patients with known oesophageal reflux disorders.

Venereal disease When treating venereal disease where co-existent syphilis is suspected, proper diagnostic procedures, including dark-field examinations, should be utilised. In all such cases monthly serological tests should be made for at least four months.

Beta-haemolytic streptococci infections Infections due to a group A beta-haemolytic streptococci should be treated for at least 10 days.

Jarisch-Herxheimer reaction Some patients with spirochete infections may experience a Jarisch-Herxheimer reaction shortly after doxycycline treatment is started. Patients should be reassured that this is a usually self-limiting consequence of antibiotic treatment of spirochete infections.

Myasthenia gravis Due to a potential for weak neuromuscular blockade, care should be taken in administering tetracyclines to patients with myasthenia gravis.

Vibramycin capsules contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

There have been reports of prolonged prothrombin time in patients taking warfarin and doxycycline. Because the tetracyclines have been shown to depress plasma prothrombin

activity, patients who are on anticoagulant therapy may require downward adjustment of their anticoagulant dosage.

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving Vibramycin in conjunction with penicillin.

The absorption of doxycycline is impaired by concurrently administered antacids containing aluminium, calcium, magnesium or other drugs containing these cations; oral zinc, iron salts or bismuth preparations.

The serum half-life of doxycycline is shortened when patients are concurrently receiving alcohol, barbiturates, carbamazepine or phenytoin.

A few cases of pregnancy or breakthrough bleeding have been attributed to the concurrent use of tetracyclines with oral contraceptives.

The concurrent use of tetracyclines and methoxyflurane has been reported to result in fatal renal toxicity.

Laboratory Test Interactions

False elevations of urinary catecholamine levels may occur due to interference with the fluorescence test.

4.6 Fertility, pregnancy and lactation

Pregnancy

Vibramycin has not been studied in pregnant patients. It should not be used in pregnancy unless, in the judgement of the physician, it is essential for the welfare of the patient. (see section 4.3 about use during tooth development).

Results of animal studies indicate that tetracyclines cross the placenta, are found in foetal tissues and can have toxic effects on the developing foetus (often related to retardation of skeletal development). Evidence of embryotoxicity has also been noted in animals treated early in pregnancy.

Breastfeeding

Tetracyclines are present in the milk of lactating women who are taking a drug of this kind and should therefore not be used in nursing mothers (see section 4.3 about use during tooth development).

4.7 Effects on ability to drive and use machines

The effect of doxycycline on the ability to drive and operate heavy machinery has not been studied. There is no evidence to suggest that doxycycline may affect these abilities.

4.8 Undesirable effects

The following adverse reactions have been observed in patients receiving tetracyclines, including doxycycline.

System Organ Class	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Not known (frequency cannot be estimated from the available data)
Blood and lymphatic system disorders			Haemolytic anaemia, neutropenia, thrombocytopenia, eosinophilia	
Immune system disorders	Hypersensitivity (including anaphylactic shock, anaphylactic reaction, anaphylactoid reaction, angioedema, exacerbation of systemic lupus erythematosus, pericarditis, serum sickness, Henoch-Schonlein purpura, hypotension, dyspnoea, tachycardia, peripheral oedema and urticaria)		Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), Jarisch-Herxheimer reaction ^b (see section 4.4)	
Endocrine disorders			Brown-black microscopic discolouration of thyroid glands	
Metabolism and nutrition disorders			Decreased appetite	
Nervous system disorders	Headache		Benign intracranial hyperthension (pseudotumor cerebri), fontanelle bulging	
Ear and labyrinth disorders			Tinnitus	
Vascular disorders			Flushing	

System Organ Class	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Not known (frequency cannot be estimated from the available data)
Gastrointestinal disorders	Nausea/vomiting	Dyspepsia (Heartburn/gastritis)	Pancreatitis, pseudomembranous colitis, <i>Clostridium difficile</i> colitis, oesophageal ulcer, oesophagitis, enterocolitis, inflammatory lesions (with monilial overgrowth) in the anogenital region, dysphagia, abdominal pain, diarrhoea, glossitis	Tooth discolouration ^a
Hepatobiliary disorders			Hepatotoxicity, hepatitis, hepatic function abnormal	
Skin and subcutaneous tissue disorders	Photosensitivity reaction, rash including maculopapular and erythematous rashes		Toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme, dermatitis exfoliative, photoonycholysis, skin hyperpigmentation ^c	
Musculoskeletal and connective tissue disorders			Arthralgia, myalgia	
Renal and urinary disorders			Blood urea increased	

CIOMS III categories: Common ≥1/100 to <1/10 (≥1% and <10%), Uncommon ≥1/1,000 to <1/100 (≥0.1% and <1%), Rare ≥1/10,000 to <1/1,000 (≥0.01% and <0.1%), Not known (frequency cannot be estimated from the available data)

b in the setting of spirochete infections treated with doxycycline.

c with chronic use of doxycycline.

a. Reversible and superficial discolouration of permanent teeth has been reported with the use of doxycycline but frequency cannot be estimated from available data

System Organ Class	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Not known (frequency cannot be estimated from the available data)
---------------------------	--------------------------------------	---	--	--

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

Acute overdosage with antibiotics is rare. In the event of overdosage discontinue medication. Gastric lavage plus appropriate supportive treatment is indicated. Dialysis does not alter serum half-life and thus would not be of benefit in treating cases of overdosage.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Tetracyclines, ATC Code: J01 AA02.

Vibramycin is primarily bacteriostatic and is believed to exert its antimicrobial effect by the inhibition of protein synthesis. Vibramycin is active against a wide range of Gram-positive and Gram-negative bacteria and certain other micro-organisms.

Vibramycin has a high degree of lipid solubility and a low affinity for calcium. It is highly stable in normal human serum. Vibramycin will not degrade into an epianhydro form.

5.2 Pharmacokinetic properties

Tetracyclines are readily absorbed and are bound to plasma proteins in varying degrees. They are concentrated by the liver in the bile and excreted in the urine and faeces at high concentrations and in a biologically active form. Doxycycline is virtually completely absorbed after oral administration. Studies reported to date indicate that the absorption of doxycycline, unlike certain other tetracyclines, is not notably influenced by the ingestion of food or milk.

Following a 200 mg dose, normal adult volunteers averaged peak serum levels of 2.6 micrograms/ml of doxycycline at 2 hours decreasing to 1.45 micrograms/ml at 24 hours. Studies have shown no significant difference in serum half-life of Vibramycin (range 18 to 22 hours) in individuals with normal or severely impaired renal function.

Haemodialysis does not alter the serum half-life of Vibramycin.

5.3. Preclinical safety data

Not applicable.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Maize starch

Lactose monohydrate

Alginic acid

Magnesium stearate

Sodium laurilsulfate

Capsule shell:

Gelatin

Titanium dioxide (E171)

Indigo carmine (E132)

Yellow iron oxide

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

4 years.

6.4 Special precautions for storage

Store below 25°C.

6.5 Nature and contents of container

Packs of 12 capsules. Aluminium/PVC blister strips, a single strip of 12 capsules in a carton box.

6.6 Special precautions for disposal and other handling

No special requirements.

7. MARKETING AUTHORISATION HOLDER

Pfizer Healthcare Ireland
9 Riverwalk,
National Digital Park
Citywest Business Campus
Dublin 24

Ireland

8. MARKETING AUTHORISATION NUMBER

PA 0822/193/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 14 October 1977

Date of last renewal: 14 October 2007

10. DATE OF REVISION OF THE TEXT

06/2019

Ref: VM 22_1